




Chinese Pharmaceutical Association
Institute of Materia Medica, Chinese Academy of Medical Sciences

Acta Pharmaceutica Sinica B

www.elsevier.com/locate/apsb
www.sciencedirect.com



REVIEW

Cytochrome P450s in algae: Bioactive natural product biosynthesis and light-driven bioproduction



Shanmin Zheng^{a,b,†}, Jiawei Guo^{b,†}, Fangyuan Cheng^b,
Zhengquan Gao^c, Lei Du^b, Chunxiao Meng^{a,*}, Shengying Li^{b,d,*},
Xingwang Zhang^{b,*}

^aSchool of Life Sciences, Shandong University of Technology, Zibo 255000, China

^bState Key Laboratory of Microbial Technology, Shandong University, Qingdao 266237, China

^cSchool of Pharmacy, Binzhou Medical University, Yantai 264003, China

^dLaboratory for Marine Biology and Biotechnology, Qingdao National Laboratory for Marine Science and Technology, Qingdao 266237, China

Received 8 November 2021; received in revised form 5 January 2022; accepted 17 January 2022

KEY WORDS

Natural product biosynthesis;
Algal pharmaceuticals;
Cytochrome P450 enzymes;
Algae;
Light-driven bioproduction

Abstract Algae are a large group of photosynthetic organisms responsible for approximately half of the earth's total photosynthesis. In addition to their fundamental ecological roles as oxygen producers and as the food base for almost all aquatic life, algae are also a rich source of bioactive natural products, including several clinical drugs. Cytochrome P450 enzymes (P450s) are a superfamily of biocatalysts that are extensively involved in natural product biosynthesis by mediating various types of reactions. In the post-genome era, a growing number of P450 genes have been discovered from algae, indicating their important roles in algal life-cycle. However, the functional studies of algal P450s remain limited. Benefitting from the recent technical advances in algae cultivation and genetic manipulation, the researches on P450s in algal natural product biosynthesis have been approaching to a new stage. Moreover, some photoautotrophic algae have been developed into “photo-bioreactors” for heterologous P450s to produce high-value added pharmaceuticals and chemicals in a carbon-neutral or carbon-negative manner. Here, we comprehensively review these advances of P450 studies in algae from 2000 to 2021.

*Corresponding authors. Tel./fax: +86 532 58632496.

E-mail addresses: mengchunxiao@126.com (Chunxiao Meng), lishengying@sdu.edu.cn (Shengying Li), zhangxingwang@sdu.edu.cn (Xingwang Zhang).

†These authors made equal contributions to this work.

Peer review under responsibility of Chinese Pharmaceutical Association and Institute of Materia Medica, Chinese Academy of Medical Sciences.

<https://doi.org/10.1016/j.apsb.2022.01.013>

2211-3835 © 2022 Chinese Pharmaceutical Association and Institute of Materia Medica, Chinese Academy of Medical Sciences. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Cytochrome P450 enzymes (P450s) are a superfamily of heme (iron protoporphyrin) proteins that are broadly distributed in human, animals, plants, algae, fungi, bacteria, archaea and even viruses^{1–4}. The name of “P450” is given owing to their distinctive spectroscopic absorption maximum at 450 nm upon its reduced heme-iron center bound with carbon monoxide⁵. Along with the explosion of genome sequencing data, a fast increasing number of P450-encoding genes, up to now more than 300,000, have been discovered from various organisms, which have greatly advanced our understanding on the important evolutionary, physiological and catalytic roles of the P450 superfamily^{6,7}. It is now well known that P450s are extensively involved in both primary and secondary metabolisms^{8–13}. For instance, many prokaryotic and eukaryotic P450s undertake essential physiological functions such as participating in the synthesis of membrane sterols, phytohormones and signaling molecules¹⁴. Meanwhile, the P450s from human or other higher animals are capable of mediating the degradation of drugs, toxins and other xenobiotics¹¹. Moreover, a tremendous number of P450s from plants and microorganisms (both bacteria and fungi) are frequently found to play essential roles in natural product biosynthesis^{3,4}.

During the biosynthesis of natural products, generally, P450s implement their functions by catalyzing the common mono-oxygenation (*e.g.*, hydroxylation and epoxidation) reactions to decorate the core structure, improve the aqueous solubility of products, and provide chemical handles for further modifications¹⁵. Intriguingly, in an increasing number of cases, P450s have been found capable of mediating diverse “uncommon” reactions for skeleton construction, including C–X (X = C/N/S) bond formation/scission, ring formation/expansion/contraction and even more vagarious reactions, which dramatically expand the chemical space of natural products^{16–20}. As a result, the P450-mediated reactions greatly increase the diversity of both the chemical structures and biological activities of natural products. Thus, the ubiquitous and versatile P450 enzymes have been recognized as one of the most significant biocatalysts in natural product biosynthesis and pharmaceutical bioproduction²¹.

Algae are a polyphyletic group of aquatic organisms that include species from the microscopic unicellular cyanobacteria to the giant kelps up to tens of meters in length^{22,23}. Most of the algal species are photosynthetic organisms, which are responsible for roughly 50% of the total photosynthesis on the earth^{5,24}. Meanwhile, algae play their fundamental ecological roles by acting as oxygen producers and the food base for almost all aquatic life. Also, upon decades of natural organic chemistry studies, algae have been recognized as a rich source of high-value bioactive natural products^{25–27}. Recent genome and transcriptome analyses have discovered numerous P450 genes involved in the basic life cycle of both cyanobacteria and eukaryotic algae. However, compared with other organisms, the knowledge on the functions of algal P450s still remains limited²⁸. Fortunately, recent technical advances in algal cultivation and genetic manipulation have led to more and more functional characterization of algal P450s^{29,30}.

Furthermore, with the rapid development of synthetic biology, the photoautotrophic algae are expected to be used as “photo-bioreactors” for heterologous P450s to produce high-value added products in a carbon-neutral or carbon-negative manner³¹. Specifically, for the reducing power-driven P450 reactions, in theory, the algal photosystem derived electrons can be directly employed to drive their catalytic cycle, thus circumventing the requirement of the complex redox partner systems and the precious reducing force NAD(P)H. Such solar-energy-driven P450-reaction systems have been successfully constructed in several algal species, in which the biosynthesis of target compounds can be fueled by the excess excitation energy during photosynthesis.

Herein, we will not only comprehensively review the important advances of algal P450 studies on natural product biosynthesis and photo-driven P450 reactions designed in algae, but also analyze the current questions/challenges and the future prospects of P450 studies in algae, aiming to inspire more cost-effective, eco-friendly and sustainable algae-based bioproduction of chemicals and pharmaceuticals in the future.

2. Algal P450 catalytic processes

Generally, the P450-mediated mono-oxygenation reactions follow a common equation of $R-H + O_2 + 2H^+ + 2e^- \rightarrow R-OH + H_2O$ ³². To initiate the reaction, two electrons originated from NAD(P)H need to be delivered to the heme-iron reactive center of P450 with the aid of redox partner protein(s) (Fig. 1)³³. Therefore, according to the importance of redox partner(s), the algal P450 catalytic systems can be classified into two major types (types I and II; Fig. 1). The prokaryotic algae (*i.e.*, cyanobacteria) adopt the three-component type I P450 catalytic system, in which the redox partners include a FAD-containing ferredoxin reductase (FdR) and a small iron-sulfur ([Fe–S]) protein ferredoxin (Fdx). By contrast, the eukaryotic algae mainly possess the Type II P450 catalytic system, wherein the P450 enzyme employs a single FAD/FMN-containing flavoprotein, referred to as cytochrome P450 reductase (CPR), to transfer the reducing equivalents. Compared with the Type I system that all the elements are cytosolically soluble proteins, the type II algal P450s and their cognate CPRs are usually endomembrane-associated proteins (Fig. 1)³⁴. This difference has led to thriving functional studies of prokaryotic P450s, while greatly limited the corresponding researches on higher algal P450s due to the technical difficulties associated with the heterologous overexpression of membrane-bound proteins³⁵.

Thereafter, the P450 sequentially accepts the two electrons shuttled from NAD(P)H by redox partner(s) to activate the inert O_2 at the heme-iron reactive center and inserts a single “O” atom to the bound substrate (R–H) to yield R–OH (Fig. 1). The detailed P450 catalytic cycle has been described in many recent reviews^{3,4,12,36}. Briefly, the intact catalytic process could be divided into two halves: in the first half cycle, the substrate-bound heme-iron center accepts two electrons, a O_2 molecule, and a pair of protons to generate the highly reactive species compound I (CpdI); and in the second half cycle, CpdI triggers the important “hydrogen-abstraction” process against the substrate R–H to




Figure 1 The two major types of algal P450s (left) and their catalytic systems which share a common catalytic cycle (right). The electron transport pathways: as shown in the left panel, NADPH first transfers a pair of electrons in the form of hydrogen ion (H^+) to FAD in either FdR or the FAD domain of CPR; then FAD transfers the two electrons one at a time to the iron-sulfur cluster ($[Fe-S]$) of Fdx or to the FMN domain of CPR; and finally Fdx or the FMN domain delivers the electron to the heme-iron center of P450 to drive the catalytic cycle as shown in the right panel³³.

produce the substrate radical (R^{\cdot}) and CpdII; then, the “OH rebound” reaction occurs to quench the R^{\cdot} and generate the monooxygenation product $R-OH$ (Fig. 1). It is worth noting that for many P450s, the catalytic cycle can be shortcut to the peroxide shunt pathway when using H_2O_2 as the sole oxygen and electron donor, in which H_2O_2 directly activates the resting heme-Fe(III) to Cpd0 (Fig. 1)^{34,37}. In rare cases, such as the fatty acid-peroxide ($R-OOH$) metabolizing P450 family members including CYP5, CYP8A, and CYP74 enzymes, the ferric enzyme can directly cleave the peroxy bond, leading to the formation of CpdII and the fatty acid alkoxyl radical (RO^{\cdot}), and finally yields the epoxyalcohol or allene oxide products (Fig. 1)³⁸.

3. Cyanobacterial P450s in natural product biosynthesis

Cyanobacteria, also referred to as blue-green algae, are a class of prokaryotic algae ubiquitously distributed in fresh and saline waters³⁹. Cyanobacteria are thought to be the oldest photosynthetic organisms that played a central role during the evolution of life by contributing to the oxygenation of the early earth. Interestingly, they are also an abundant source of bioactive natural products, including polyketides (PKs), non-ribosomal peptides (NRPs), terpenoids and the hybrids thereof^{5,40}. Genome mining revealed that cyanobacteria represent a huge arsenal of P450s, which may function as indispensable roles in both primary and secondary metabolisms⁴¹. In this section, we will summarize the identified P450-catalyzed reactions involved in cyanobacterial natural product biosynthesis (Table 1).

3.1. All-trans-retinoic acid

All-trans-retinoic acid (**1**) is a naturally occurring vitamin A-derived pharmaceutical that is highly effective in the treatment of acute promyelocytic leukemia, and it shows the potential to treat some autoimmune diseases, such as psoriasis, systemic lupus erythematosus, inflammatory arthritis, and ulcerative colitis^{42,43}. Genome mining of *Synechocystis* sp. PCC 6803 revealed a function unknown P450 named CYP120A1⁴⁴. In an earlier study, CYP120A1 was proposed to participate in the transformation of **1**

based on bioinformatics analysis and the co-crystal structure with **1**⁴⁵. Later, an *in vitro* bioassay, using the microsomal fraction of the recombinant *Saccharomyces cerevisiae* WAT11 expressing CYP120A1, showed that CYP120A1 is indeed capable of hydroxylating C16 and C17 of **1** to form **2** and **3** respectively (Fig. 2A)⁴⁶. It is noteworthy that CYP120A1 is the first identified retinoic acid-metabolizing enzyme of non-animal origin⁴⁵.

3.2. Type A malyngamides

Type A malyngamides are a family of marine cyanobacteria derived natural products that have potent anticancer and anti-inflammatory activities⁴⁷. Recently, malyngamide I (**4**) and malyngamide C (**5**) were isolated from the cyanobacterial strains *Okeania hirsute* PAB and PAP respectively, and their biosynthetic pathways were characterized based on *in vivo* studies. On the early stage, the carbon skeleton is assembled by a polyketide synthase (PKS) and a nonribosomal peptide synthetase (NRPS) to form **6** and **7** with the aid of several tailoring enzymes. Next, P450 MgcT catalyzes the $C4=C9$ oxidation of **6** to produce the epoxidation product, which is further hydroxylated at the C8 position by another P450 MgcU to form the precursor of **4** in the strain PAB (Fig. 2B). Similarly, in *O. hirsute* PAP, the MgcT counterpart P450 MgiT was considered responsible for the $C4=C9$ epoxidation of **7** to produce **5** (Fig. 2B)⁴⁸.

3.3. Cryptophycins

Cryptophycins, as exemplified by cryptophycin 2 (**8**), are a class of peptolide natural products isolated from the lichen cyanobacterial symbiont *Nostoc* sp. ATCC 53789 and *Nostoc* sp. GSV224⁴⁹. They are potent anticancer agents by stimulating cellular microtubule instability, inhibiting microtubule assembly, and inducing tubulin self-association⁵⁰. In the biosynthetic study of **8**, a P450 enzyme CrpE was characterized to mediate the formation of the characteristic β -epoxy ring at the $C2=C3$ double bond towards the substrate cryptophycin 4 (**9**) (Fig. 2C), based on the *in vitro* enzymatic assay using the purified CrpE protein expressed in the recombinant *Escherichia coli*⁵¹. In addition, CrpE

Table 1 The functions of algal P450s.

P450	Biosynthetic pathway	Function	<i>In vitro/in vivo</i> characterization	Source	Prokaryotic (P)/ Eukaryotic (E)	Ref.
AB2	Polychlorinated biphenyls/triphenyls	C–C or C–O biaryl ring coupling	Y/Y	<i>F. ambigua</i> 108b	P	60
AB3	Polychlorinated biphenyls/triphenyls	C–C or C–O biaryl ring coupling	Y/Y	<i>F. ambigua</i> 108b	P	60
CrpE	Cryptophycin 2	Epoxidation	Y/–	<i>Nostoc</i> sp. ATCC 53789 and <i>Nostoc</i> sp. GSV224	P	51
CYP120A1	All- <i>trans</i> -retinoic acid	Hydroxylation	Y/–	<i>Synechocystis</i> sp. PCC 6803	P	44–46
CYP110	Long-chain saturated fatty acids	Hydroxylation	–/–	<i>Anabaena</i> sp. PCC 7120	P	62,63
<i>Cke</i> CYP97A1	Xanthophylls	Hydroxylation	–/Y	<i>Chlorella kessleri</i>	E	112
<i>Db</i> CYP97A	Xanthophylls	Hydroxylation	–/Y	<i>Dunaliella bardawil</i>	E	80
<i>Db</i> CYP97B	Xanthophylls	Hydroxylation	–/Y	<i>D. bardawil</i>	E	80
<i>Db</i> CYP97C	Xanthophylls	Hydroxylation	–/Y	<i>D. bardawil</i>	E	80
ESEAS	Epoxyalcohols	Epoxidation	Y/–	<i>Ectocarpus siliculosus</i>	E	75
<i>Eg</i> CYP97H1	Xanthophylls	Hydroxylation	–/Y	<i>Euglena gracilis</i>	E	81
<i>Hae</i> CYP97A	Xanthophylls	Hydroxylation	–/Y	<i>Haematococcus pluvialis</i>	E	78
<i>Hae</i> CYP97B	Xanthophylls	Hydroxylation	–/Y	<i>H. pluvialis</i>	E	78
<i>Hae</i> CYP97C	Xanthophylls	Hydroxylation	–/Y	<i>H. pluvialis</i>	E	78
HctG	Hectochlorin	Hydroxylation	–/–	<i>Lyngbya majuscula</i>	P	53
HctH	Hectochlorin	Hydroxylation	–/–	<i>L. majuscula</i>	P	53
KfAOS	Jasmonates	Epoxidation	Y/–	<i>Klebsormidium flaccidum</i>	E	74
LtxB	Lyngbyatoxin A	C–N bond formation	Y/–	<i>Lyngbya majuscula</i>	P	66
MgcT	Malylgamide C	Epoxidation	–/Y	<i>Okeania hirsute</i> strains PAB	P	48
MgcU	Malylgamide C	Hydroxylation	–/Y	<i>O. hirsute</i> strains PAB	P	48
MgiT	Malylgamide I	Epoxidation	–/Y	<i>O. hirsute</i> strains PAP	P	48
P450NS	Oxidized germacrene A	Oxidative cyclization	–/Y	<i>Nostoc</i> sp. PCC 7120	P	57,58
<i>Pu</i> CHY1	Xanthophylls	Hydroxylation	–/Y	<i>Porphyra umbilicalis</i>	E	113
<i>Ptr</i> CYP97B1	Xanthophylls	Hydroxylation	–/Y	<i>Phaeodactylum tricorutum</i>	E	114
<i>Ptr</i> CYP97B2	Xanthophylls	Hydroxylation	–/Y	<i>P. tricorutum</i>	E	114

can also convert the **9**-analogues, **10**–**13**, to the corresponding epoxidation products **14**–**17** *in vitro* (Fig. 2C), indicative of a broad substrate spectrum⁵².

3.4. Hectochlorin

Hectochlorin (**18**) is another peptolide product identified from marine cyanobacteria *Lyngbya majuscula* and shows remarkable antifungal and cytotoxic properties. Biosynthetically, **18** is produced by a PKS-NRPS hybrid assembly-line, during which the elongating substrates are tethered to a peptidyl carrier protein (PCP)⁵³. A few P450s have been reported to be capable of mediating the online oxidations, of which the P450's substrate is linked to an acyl carrier protein (ACP) or PCP, thus involving the protein–protein interactions during the substrate recognition process. The biosynthetic gene cluster of **18** encodes two P450s HctG and HctH⁵³. According to bioinformatics analysis and the biocatalytic principles, HctG/HctH were proposed to catalyze such an online hydroxylation at the C3 position of the PCP-tethered substrate **19** to form **20**, prior to the NPRS-mediated cyclopeptide formation (Fig. 2D)⁵³.

3.5. Fatty acids

The ω -hydroxylated fatty acids are important starting materials in chemical, pharmaceutical, and cosmetic industries⁵⁴. However, the regiospecific ω -hydroxylation of the inert terminal sp^3 C–H bond in a hydrocarbon chain is thermodynamically disfavored for chemical synthesis. Some P450s have long been found to hold the

capacity of mediating such ω -hydroxylation reactions towards various fatty acid substrates^{55,56}. Genome mining of the freshwater cyanobacterium *Anabaena* sp. PCC 7120 revealed an unknown P450 named CYP110^{57,58}. Protein sequence analysis indicated the high similarity of CYP110 with the mammalian CYP4 family proteins and the bacterial P450 BM3 from *Bacillus megatarium*, two classes of identified fatty acid ω -hydroxylases (Fig. 2E). The substrate titration experiments and spectroscopic spin-shift assays demonstrated that the saturated and unsaturated fatty acids of chain lengths C₁₀–C₂₀ could bind with CYP110⁵⁸. Thus, CYP110 was proposed to catalyze ω -hydroxylation of long-chain fatty acid (**21**) to form **22** (Fig. 2E)^{57,58}.

3.6. Polyhalogenated aromatics

Apart from the common monooxygenation reactions, P450s can also mediate various complex structure-reshaping reactions. Enzymatic aromatic/phenolic couplings are fascinating reactions in natural product biosynthesis, because these regio- and/or stereoselective transformations are highly challenging in chemical synthesis. Polyhalogenated aromatics represent a class of anthropogenic chemicals that are widely used in agriculture⁵⁹. Intriguingly, this class of molecules have also been isolated from some marine microorganisms, such as the *Fischerella ambigua* 108b originated polychlorinated triphenyl product **23** and biphenyl products **24** and **25**⁶⁰. The biosynthetic gene cluster of **23**–**25** was recently revealed by genome mining, which is comprised of ten genes including two P450 genes (*AB2* and *AB3*). Based on *in vivo* precursor feeding experiments, *AB2* and *AB3* were elucidated to




Figure 2 The monooxygenation reactions catalyzed by cyanobacterial P450s. (A) The all-*trans*-retinoic acid (**1**) hydroxylation reactions catalyzed by CYP120A1 from *Synechocystis* sp. PCC 6803. (B) The epoxidation and hydroxylation reactions catalyzed by different P450s during the biosynthesis of malyngamides **4** and **5**. (C) The CrpE-catalyzed epoxidation reactions of cryptophycin 4 (**9**) and **9** analogues (**10–13**). (D) The online hydroxylation catalyzed by P450 HctG/HctH during the biosynthesis of **18**. (E) The ω -hydroxylation of fatty acid catalyzed by CYP110.

catalyze the C3–C3' or C3–O1' coupling of the monomer **26** to form **24** and **25**, and the O1–C4'/C6'–C3'' coupling of **26** and **27** to yield **23** (Fig. 3A)⁶⁰. The *in vitro* reconstituted enzymatic reactions disclosed more catalytic details, of which AB2 and AB3 can both catalyze the formation of **24** and **25**, but AB2 tends to produce **24** while AB3 prefers the production of **25**⁶⁰. Moreover, the co-incubation of AB2 and AB3 with the monomers **26** and **27** generated the tri-phenol product **23** (Fig. 3A)⁶⁰.

Mechanistically, the P450-catalyzed aromatic/phenolic coupling reactions are considered to undergo a biradical coupling pathway. In this case, AB2/AB3 primarily abstracts a single hydrogen from the phenolic hydroxyl group of **26** or **27** to yield the oxygen radicals **i** or **ii**, which can undergo the electron rearrangement to give the corresponding carbon radical **iii** and **iv**. Then, two molecules of **iii** couple to yield **24**, while **i** and **iii** couple to produce **25**. As for the formation of **23**, after the generation of **v** from **i** and **iv**, AB2/AB3 further abstracts the H6' to give a new radical **vi**, which couples with **iii** to yield the tri-phenol products (Fig. 3A)⁶¹.

3.7. Germacrene A

Germacrene A (**28**), derived from the precursor of farnesyl pyrophosphate (**29**), is a general biosynthetic intermediate of many

bioactive sesquiterpenes, such as englerin A and guaianolide, in both plants and microorganisms⁶². In a genome mining study of the cyanobacterium *Nostoc* sp. PCC 7120, a sesquiterpene synthase (NS1) was found to be able to biosynthesize **28**⁶³. Stepwise heterologous expression of NS1 and the co-clustered P450NS in *E. coli* revealed that P450NS could mediate an uncommon oxidative cyclization of **28** to yield the more hydrophilic product **30** (Fig. 3B)⁶⁴. It was proposed that P450NS first epoxidizes the C5=C6 double bond to produce the unstable intermediate **i**, which could undergo C–O bond heterolysis to give **ii**. Eventually, **30** is generated after intramolecular electron rearrangement and ring-closure of **ii** (Fig. 3B).

3.8. Lyngbyatoxin A

Lyngbyatoxin A (**31**) is an indole alkaloid cyanotoxin that can activate the protein kinase C (PKC) isozyme, and also a potent irritant and vesicant, as well as a carcinogen⁶⁵. Biogenetically, the C–N bond formation is of special interest since such a C–H amination reaction is infrequently occurring in nature. The biosynthetic gene cluster of **31** was mined from a Hawaiian cyanobacterium strain *Lyngbya majuscula*, which contains a P450 encoding gene *ltxB*. Based on the *in vitro* reconstituted enzymatic reaction, LtxB was determined to mediate the conversion of the




Figure 3 The uncommon reactions catalyzed by cyanobacterial P450s and the related catalytic mechanisms (in box). (A) The C–C and C–O coupling reactions catalyzed by AB2 and AB3 during the biogenesis of polyhalogenated aromatics. (B) The oxidative cyclization reaction mediated by P450NS. (C) The C–N bond formation catalyzed by LtxB in the biosynthesis of lyngbyatoxin A (**31**), and the various C–N bond bearing products (**34**–**38**) formed by LtxB. (D) The C–S bond formation catalyzed by TleB.

substrate **32** to **33** by catalyzing the rare C4–N13 bond formation reaction. To explore the substrate tolerance of LtxB, several analogues of **32** were synthesized and reacted with LtxB *in vitro*. Intriguingly, all these analogues could be converted into the corresponding C–N coupling products (**34**–**38**, Fig. 3C)⁶⁶. Recently, the mechanistic study of TleB, a homologous protein of LtxB from *Streptomyces*, was conducted by crystal structure analysis, revealing a biradical-coupling process (route 1, Fig. 3C) or an alternative carbocation involved nucleophilic pathway (route 2, Fig. 3C)⁶⁵. Moreover, in a following study, it was revealed that TleB could also catalyze a C–S bond formation towards the substrate *S*-analogue **39** to yield **40** (Fig. 3D)²⁰. These results together prove LtxB/TleB to be potential bio-catalysts for generation of the C–N/C–S bond containing products.

4. Eukaryotic algal P450s in natural product biosynthesis

Eukaryotic algae encompass diverse photosynthetic species ranging from unicellular microalgae to multicellular giant kelps^{67,68}. Continuous natural product prospecting has revealed that eukaryotic algae are also an important source of bioactive

compounds. The most attractive merit is that some eukaryotic algae can produce abundant bioactive carotenoid-like pigments and halogenated natural products^{69,70}. A recent genome survey disclosed the high diversity of P450s in eukaryotic algae²⁸. However, due to the technical limitations in laboratory cultivation and gene-manipulation of eukaryotic algae, the progress on these membrane-bound P450s remains limited, especially for the P450s from giant kelps^{25,27}. In this part, a limited number of eukaryotic algal P450s and their catalytic reactions will be reviewed (Table 1).

4.1. Jasmonates

Jasmonates, taking the 12-oxo-phytodienoic acid (**41**) and jasmonic acid (**42**) as examples, are important hormones distributed in plants and algae. They are also potential therapeutic agents for several types of cancer⁷¹. During the biosynthesis of **41** and **42**, the fatty acid-derived peroxide oxylipins, such as 13-hydroperoxy-(*E,Z,Z*)-9,11,15-octadecatrienoic acid (**43**), are the key intermediates⁷². The P450 KfAOS was identified as a CYP74 family member involved in jasmonate biosynthesis from the green microalgae *Klebsormidium flaccidum*. Functional characterization revealed that KfAOS is able to convert **43** into the oxylipin allene

oxide (**44**, Fig. 4A)³⁸. Then, the unstable product **44** undergoes a spontaneous rearrangement to form **41**, and such a transformation can be accelerated by an allene oxide cyclase (Fig. 4A)^{73,74}. In another study, the CYP74-related P450 ESEAS (CYP5164B1) from the brown algae *Ectocarpus siliculosus* was found to catalyze a similar reaction⁷⁵. As reported, ESEAS is capable of transforming both (9*S*,10*E*,12*Z*)-9-hydro (peroxy)-10,12-octadecadienoic acid (**45**) and (9*Z*,11*E*,13*S*)-13-hydro (peroxy)-9,11-octadecadienoic acid (**46**) into the corresponding epoxyalcohol products **47** and **48**, respectively (Fig. 4B).

Intriguingly, neither redox partners nor electron donor is required for these CYP74-catalyzed reactions. As for the catalytic mechanism, it is proposed that after binding with the fatty acid hydroperoxide (**49**), the heme-Fe(III) of CYP74 (KfAOS or ESEAS) homolytically cleaves the O–O bond to give a [RO[•]] radical (**50**) and CpdII (Fe(IV)–OH). Then, the [RO[•]] radical

reacts with the adjacent double bond to form the epoxy species and relocates the radical to the α -carbon, forming the carbon radical **51**. Thereafter, the [HO[•]] of CpdII rebounds to **51** to yield the epoxyalcohol product **52** (for ESEAS). Alternatively, CpdII further abstracts a hydrogen from the α -carbon of **51** to afford the allene oxide product **53** and release a water molecule (for KfAOS) (Fig. 4C)^{38,75,76}. Thus, this is a unique class of P450s to catalyze unusual isomerization reactions instead of the normal oxygenation reactions, because no additional electron and oxygen is introduced into the catalytic cycle.

4.2. Xanthophylls

Xanthophylls, such as lutein (**54**), are a large class of oxygen-containing carotenoid pigments that play important protecting roles in photosynthesis of eukaryotic algae and higher plants^{77,78}.




Figure 4 The reactions catalyzed by eukaryotic algal P450s and the related enzymatic mechanism (in box). (A) The oxylipin allene oxide (**44**) epoxidation catalyzed by KfAOS from the green microalgae *Klebsormidium flaccidum* during the formation of **41** and **42**. (B) The epoxidation reactions catalyzed by ESEAS from the brown algae *Ectocarpus siliculosus*. (C) The putative mechanisms of KfAOS and ESEAS. (D) The hydroxylation reactions of α -carotene (**55**), α -cryptoxanthin (**56**) and zeinoxanthin (**57**) catalyzed by DbCYP97C from the green algae *Dunaliella bardawil* during the biosynthesis of lutein (**54**). (E) The hydroxylation reaction of β -cryptoxanthin (**59**) catalyzed by EgCYP97H1 from *Euglena gracilis*. (F) The CYP51 mediated 14 α -demethylation reaction during sterol biosynthesis.

Physiologically, **54** and its structural analogues can absorb excess light energy to prevent photo-damages to host organisms. Xanthophylls also have health benefits and therapeutic potentials against neurologic, ophthalmologic, oral, allergic and immune diseases. Biosynthetically, **54** is derived from α -carotene (**55**) with two hydroxylation steps⁷⁹. A recent study on carotene hydroxylases from the green algae *Dunaliella bardawil* revealed a group of P450s (DbCYP97A, DbCYP97B, and DbCYP97C) are responsible for the C4/C5' hydroxylation of **55** to form **54** (Fig. 4D). According to the functional complementation experiments in *E. coli*, the catalytic preferences of the three P450s were revealed: DbCYP97A and DbCYP97C showed a high catalytic activity toward the β -ring and ϵ -ring of **55** to yield α -cryptoxanthin (**56**) and zeinoxanthin (**57**), respectively; while DbCYP97B displayed minor hydroxylation activity toward the β -ring of **55** (Fig. 4D)⁸⁰. Interestingly, unlike other eukaryotic P450s, the CYP97 proteins lack a transmembrane helix, thus being peripherally bound to the membrane. In addition, another CYP97 member *EgCYP97H1* from *Euglena gracilis* was characterized as a β -carotene monohydroxylase, which hydroxylates the C4 position of β -carotene (**58**) to form β -cryptoxanthin (**59**, Fig. 4E). Notably, this is the first example that a P450 functions as a β -carotene hydroxylase⁸¹.

4.3. Sterols

Sterols are important structural and regulatory components in eukaryotes⁸². As observed in plants, CYP51 is a necessary P450 for sterol biosynthesis by catalyzing the 14 α -demethylation of obtusifolium (**60**) to yield **61**, via the putative geminal diol and carbocation intermediates (Fig. 4F)⁸³. However, the algal CYP51 had not been investigated until a recent study carried out in *Nannochloropsis oceanica*. As reported, addition of the CYP51 inhibitor tebuconazole (TEB) into the culture of *N. oceanica* led to the accumulation of the major biosynthetic intermediate of sterols⁸⁴. Accordingly, CYP51 in *N. oceanica* was proposed to play a similar role as their homologous P450s in other eukaryotic organisms.

In summary, as for the algal P450-mediated biosynthesis of essential metabolites, analogous P450-catalyzed reactions can also be found in other non-algal organisms. For example, the all-*trans*-retinoic acid (**1**) metabolizing reaction can be accomplished by both the cyanobacterial P450 CYP120A1 and some human P450s (Fig. 2A)⁸⁵; the C–N bond formation mediated by cyanobacterial P450 LtxB also occurs in *Streptomyces* (Fig. 3C)⁸⁶; and the uncommon rearrangement reactions catalyzed by the eukaryotic algal CYP74 family member have also been elucidated in other higher plants (Fig. 4A–C)⁸⁷. These cross-species functional similarities may suggest some evolutionary relationship between algal and non-algal organisms. By contrast, there also exist some algae-specific reactions that have not been observed in other organisms, such as the tripolymer production co-mediated by the cyanobacterial P450s AB2 and AB3 (Fig. 3A). Besides, algae have evolved many unique P450s giving rise to the specialized natural products of algal characteristics, as exemplified by **4**, **5**, **8**, and **18**. With no doubt, these algal P450s significantly expand the catalytic diversity of the whole P450 superfamily.

5. Algae as photo-bioreactors of heterologous P450s

Algae are photoautotrophic organisms that can use the solar energy and recycle CO₂ to synthesize primary and secondary metabolites (Fig. 5)³¹, but the solar-to-biomass energy conversion

rate is quite low with the energy converting efficiencies only ranging from 1% to 2%⁸⁸. In fact, in the photosynthetic pathway, the photosystem II can efficiently absorb the solar energy for photolysis of water molecule to generate electrons for downstream metabolic pathways. However, the total cellular biosynthetic electron-demand is far less than the produced electrons^{88,89}. Therefore, theoretically, these excessively absorbed solar energy can be further utilized by introducing heterogenous catalytic elements provided that they can interact with the photosystem proteins to acquire electrons.

With regard to P450s, considering the facts that 1) the P450-catalyzed reactions need to consume a lot of NAD(P)H to gain the reducing power (electrons) to drive the catalytic cycle (Fig. 1); 2) excess reducing power is generated during algal photosynthesis⁹⁰; and 3) P450s can use the photosynthesis-associated Fdxs to shuttle the required electrons (Fig. 5), a methodology which can channel the excess reducing power from the algal photosystem to the P450 catalytic system would be highly valuable for heterologous P450s to produce high-value products by directly using the solar energy⁹¹. Technically, a heterologous P450 can be readily fused with the photosynthetic system by anchoring it into the thylakoid membrane that harbors the whole photosynthetic pathway (Fig. 5). For instance, a eukaryotic membrane-bound P450 can be directly anchored into the prokaryotic thylakoid membrane by its native membrane-associated N-terminal sequence; while the prokaryotic Type I P450 could be re-located into the thylakoid membrane by fusing it with the subunit of photosystem protein (Fig. 5). Based on this strategy, several light-driven P450 reaction systems have been successfully developed in different algal species^{31,91–96}. In this section, we will discuss the state-of-the-art algal photo-bioreactors for light-driven P450 reactions.

5.1. Caffeic acid

Caffeic acid (**62**), a phenolic natural product synthesized by all plant species, shows antioxidant, anti-inflammatory and anticancer activities and is widely used in cosmetics industry^{97–99}. At present, the commercial **62** is chemically synthesized, which is suffering from the low yield and serious environmental concerns^{91,100}. The P450 CYP98A3 (C3H) from *Arabidopsis thaliana* was identified as a caffeic acid synthase by hydroxylating the C3 position of *p*-coumaric acid (**63**). Previously, functional expression of C3H failed in a bacterial host due to its instability, low abundance, and membrane-bound nature⁹¹. In a pilot study, the possibility of using cyanobacterium as a “photo-bioreactor” to produce **62** was explored. When the codon-optimized C3H-encoding gene was introduced into *Synechocystis* sp. PCC 6803, the plant P450 was successfully expressed and located onto the thylakoid membrane⁹¹. Together with other biosynthetic enzymes, the expected product **62** was produced in the cyanobacterium under the photosynthetic growth conditions (Fig. 5). This study demonstrated the algal thylakoid to be a suitable platform for studying the membrane-bound P450s¹⁰¹.

5.2. Dhurrin

Dhurrin (**64**) is a cyanogenic glycoside produced by many plants, which could comprise up to 30% of the dry mass of etiolated sorghum seedlings¹⁰². Compound **64** exhibits strong antimicrobial properties while its cytotoxicity also discourages herbivory of some insects and animals. CYP79A1 is involved in the **64** biosynthetic pathway from *Sorghum bicolor* that converts L-




Figure 5 The algal photosynthetic pathway and its application for driving the P450-catalyzed reactions (adopted from Ref. 31 Copyright © 2021 American Chemical Society). During the photosynthesis, photosystem II (PSII) conducts the water photolysis and produces the two electrons which are then transferred to plastoquinone (PQ). PQ gains the electrons and two protons to form PQH₂, which is then oxidized by cytochrome *b*₆*f* complex (Cyt *b*₆*f*) to capture the electrons and restore PQ. Thereafter, the electrons in Cyt *b*₆*f* is stepwise transferred to Fdx via the electron-transfer protein plastocyanin (PC) and photosystem I (PSI). Then, Fdx transfers the electron to ferredoxin–NADP⁺ reductase (FNR) to produce NADPH. When the heterologous P450 is present, Fdx can direct partial electrons to P450s to drive various oxidative reactions.

tyrosine (**65**) into the N–OH bearing product **66**. When the soluble catalytic domain of CYP79A1 was fused with the cyanobacterial photosystem I derived subunit to locate it onto thylakoid, the eukaryotic P450 was successfully expressed in *Synechococcus* sp. PCC 7002; and it showed the light-dependent activity both *in vitro* and *in vivo* (Fig. 5)⁹³. During the biosynthesis of **64**, another P450 named CYP71E1 can further oxidize **66** into the nitrile-carrying compound **67**. Thus, to achieve total biosynthesis of **64**, CYP79A1 and CYP71E1 were co-expressed in *Synechocystis* sp. PCC 6803, and the results showed that the reconstituted activities of the two P450s were strictly light dependent. When the glycosyltransferase UGT85B1 was further introduced, the final product **64** was successfully produced in the cyanobacterial host (Fig. 5)⁹⁵. In another study, CYP79A1 was also successfully expressed in the eukaryotic green algae *Chlamydomonas reinhardtii*. As expected, after the CYP79A1 encoding gene was incorporated into the chloroplast genome of *C. reinhardtii*, the P450 enzyme was stably expressed and located in the chloroplast membrane⁹⁴.

5.3. Artificial chemicals

In addition to bioactive natural products, the cyanobacterial hosts have also been developed to produce high-value chemicals by running the light-driven P450 reactions. Cyclohexanol (**68**) is a key precursor in chemical/pharmaceutical industry for the production of ϵ -caprolactone, adipic acid and several polymers¹⁰³. However, **68** is intricate to acquire through chemical synthesis mainly due to the difficulty in the inert *sp*³ C–H bond activation; thus, enzymatic synthesis provides an alternative solution to address this issue¹⁰⁴. A P450 enzyme CYP-CHX from *Acidovorax* sp. CHX100 was previously identified to be such a hydroxylase that can hydroxylate cyclohexane (**69**) to **68**¹⁰⁵. To achieve the production of **68**, the P450 was expressed in the cyanobacterium *Synechocystis* sp. PCC 6803, and gram-scale of **68** was successfully produced by supplying **69** to the recombinant cyanobacterial culture (Fig. 5). Essentially, it was confirmed that the production of **68** was light dependent¹⁰⁶.

5.4. Pollutant degradation

The numerous man-made chemicals, such as the widely used pesticides and antibiotics, have greatly promoted social development in the last century. However, the environmental pollution caused by artificial chemicals also threatens the living world. The ubiquitously distributed algae have been designed as pollutant degrading-hosts by expressing biocatalysts which can mediate biodegradation reactions. Atrazine (**70**) is one of the most widely used pesticides in the world, and has been recognized as a serious environmental pollutant¹⁰⁷. Previous studies have shown that the human or mammalian P450 CYP1A1 has the capacity of converting **70** into the deisopropylated and deethylated product of atrazine (**71**), both *in vitro* and *in vivo*^{108,109}. When CYP1A1 from *Rattus norvegicus* (brown rat) was expressed in the cyanobacterium *Synechococcus* sp. PCC 7002, it was found that the protein could be located onto thylakoid membrane through its native N-terminal membrane-bound domain. Importantly, *Synechococcus* sp. PCC 7002 expressing CYP1A1 successfully degraded **70** into **71** in a light-dependent manner (Fig. 5)¹⁰⁷.

Taken together, these results demonstrate the great potential of both cyanobacteria and eukaryotic algae as light-driven P450 bioreactors and biosynthetic platforms for the production of high-value added pharmaceuticals/chemicals (or the degradation of pollutants) in a cost-effective and eco-friendly manner. Intriguingly, a recent study showed that the expression of the above-mentioned CYP1A1 in *Synechococcus elongatus* PCC 7942 led to not only an improvement of P450 activity, but also a significant increase in the quantum efficiency of photosystem⁸⁹. It was hypothesized that the introduced heterologous metabolism might provide an outlet for the excessively captured energy, thus reducing the flow to the photoprotective pathways of energy consumption. Such a balance on the absorbed light energy and metabolic capacity is physiologically important for the photosynthetic microorganism.

Despite these significant progresses, there remain a number of substantial obstacles to develop the algal “bioreactors” into “cell-factories” for industrial applications. The major problem is the low

catalytic efficiency when expressing the heterologous P450s in algae. Therefore, increasing the compatibility of the heterologous P450s with the photosystem, thus improving the electron utilizing efficiency for P450s from photosystem, would become one of the main directions of P450 photo-bioreactor improvement in the future⁹⁰. At present, some efforts have optimized the electron transfer from photosystem I to P450 by fusion-expression of P450 and Fdx^{90,110}, or through lowering the affinity of Fdx and FNR by point mutation of FNR to redirect the electrons to P450s¹¹¹. We envision that more pathbreaking strategies/approaches are required and will emerge to solve this central problem associated with the algal photo-driven P450 bioreactors or cell-factories.

6. Questions and challenges

In the post-genome era, big data analysis of algal genomes and transcriptomes has helped revealing more roles of P450s in algal living activities including both primary and secondary metabolisms. However, when comparing with other organisms, such as human, mammals, plants, fungi and bacteria, our understanding on the functions and mechanisms of algal P450s is very limited. Despite the recent significant progresses on algal P450s as summarized in this review, the breakthrough in algal P450 studies has yet to come due to some unsolved bottlenecks, especially for the eukaryotic algal P450s. The main problems associated with algal P450 functional studies lies in: 1) For cyanobacteria, genome-editing remains a time-consuming task because of its low growth rate. For some species of diploid or multiploid, the genetic manipulation is even more complicated. Besides, the expression levels of heterologous P450 genes in cyanobacteria are usually low, thus leading to inefficient synthesis of the products of interest; 2) For eukaryotic algae, genetic manipulation and heterologous P450 expression are even more difficult due to the cellular and genomic complexity, low growth rate, membrane-bound protein limitation, and unavailability of the whole genomic blueprints (especially for giant kelps); 3) For both cyanobacteria and eukaryotic algae, genome mining and biosynthetic gene cluster analysis revealed that many P450 genes are not clustered with other biosynthetic genes, making it difficult to predict and identify their catalytic functions^{28,41}; and 4) even though the current studies have shown great potential for algae as the light-driven “bioreactors” in the production of useful pharmaceuticals/chemicals, the productivity and compatibility of algal hosts for heterologous P450 expression are still too low to be applied in practical applications.

7. Conclusions and prospects

Algae are a huge group of life on the earth and play essential roles in global substance- and energy-cycles. P450s are one of the key catalytic elements for algae by mediating various common and uncommon reactions¹³. Functional and mechanistic characterization of algal P450s will provide us the knowledge on what they do and how they work in the native metabolic networks of algae. In turn, the knowledge would enable us to employ these algal P450 biocatalysts or the algal photosynthetic platform to produce natural and unnatural but useful products, such as pharmaceuticals and fine chemicals, with the accumulating methodologies and tools of synthetic biology. In recent years, more algal P450s have been functionally and mechanistically elucidated, which expands our understandings on catalytic diversity of the whole P450 superfamily. In addition, the photosynthetic algae have shown their great potential as “photo-

bioreactors” for heterogenous P450s of different origins to produce high-value products.

Finally, we conclude that P450 studies in algae are still on the early stage. Comparing with other organisms, there currently lacks a general model strain for either prokaryotic or eukaryotic algal P450 studies. However, once developed, an algal platform strain might be advantageous over *E. coli* and *Saccharomyces cerevisiae* because it would optimally have all the necessary elements, including the endomembrane system, excess O₂ supply, and different sources of reducing power, for P450 expression and functionalization. It is still questionable whether the native algal P450s can directly interact with the photosystem to channel the solar energy derived electrons to drive their catalytic cycles; and whether the Fdx-associated photosystem can be universally used as a general electron donor of P450s to replace the functionality of redox partner proteins and NAD(P)H. Taken together, we anticipate that the studies on algal P450s and “photo-bioreactors” will become more popular and feasible in the near future, along with the expansion of the synthetic biology toolbox, and the development of higher-throughput omics technologies and higher-efficiency genome-editing methods.

Acknowledgments

This work was supported by the National Key Research and Development Program of China (2020YFA0907900), the National Natural Science Foundation of China (32000039, 32025001, 31972815 and 42176124), the Natural Science Foundation of Shandong Province (ZR2019ZD20, ZR2019ZD17 and ZR2020ZD23), the Fundamental Research Funds of Shandong University (2019GN031), and the Scientific Research Fund of Binzhou Medical University (BY2021KYQD25).

Author contributions

Xingwang Zhang, Shengying Li, and Chunxiao Meng conceived the review. Shanmin Zheng, Jiawei Guo and Fangyuan Cheng summed up the literatures and prepared the manuscript. Xingwang Zhang, Shengying Li, Zhengquan Gao and Lei Du revised the manuscript. All of the authors have read and approved the final manuscript.

Conflicts of interest

The authors declare no competing financial interest.

References

1. Nelson DR. Cytochrome P450 diversity in the tree of life. *BBA-Proteins Proteom* 2018;**1866**:141–54.
2. Montellano PROD. *Cytochrome P450-structure, mechanism, and biochemistry*. 4th ed. Switzerland: Springer; 2015.
3. Zhang X, Li S. Expansion of chemical space for natural products by uncommon P450 reactions. *Nat Prod Rep* 2017;**34**:1061–89.
4. Zhang X, Guo J, Cheng F, Li S. Cytochrome P450 enzymes in fungal natural product biosynthesis. *Nat Prod Rep* 2021;**38**:1047–228.
5. Robert FO, Pandhal J, Wright PC. Exploiting cyanobacterial P450 pathways. *Curr Opin Microbiol* 2010;**13**:301–6.
6. Sherman DH, Li S, Yermalitskaya LV, Kim Y, Smith JA, Waterman MR, et al. The structural basis for substrate anchoring, active site selectivity, and product formation by P450 PikC from *Streptomyces venezuelae*. *J Biol Chem* 2006;**281**:26289–97.

7. Cryle MJ, Schlichting I. Structural insights from a P450 Carrier Protein complex reveal how specificity is achieved in the P450(BioI) ACP complex. *Proc Natl Acad Sci U S A* 2008;**105**:15696–701.
8. Kittendorf JD, Sherman DH. The methymycin/pikromycin pathway: a model for metabolic diversity in natural product biosynthesis. *Bioorg Med Chem* 2009;**17**:2137–46.
9. Murphy CD. Drug metabolism in microorganisms. *Biotechnol Lett* 2015;**37**:19–28.
10. Theis T, Backhaus T, Bossmann B, Grimme L. Xenobiotic biotransformation in unicellular green algae. Involvement of cytochrome P450 in the activation and selectivity of the pyridazinone pro-herbicide metflurazon. *Plant Physiol* 1996;**112**:361–70.
11. Wrighton SA, Stevens JC. The human hepatic cytochromes P450 involved in drug metabolism. *Crit Rev Toxicol* 1992;**22**:1–21.
12. Rudolf JD, Chang CY, Ma M, Shen B. Cytochromes P450 for natural product biosynthesis in streptomycetes: sequence, structure, and function. *Nat Prod Rep* 2017;**34**:1141–72.
13. Hansen CC, Nelson DR, Møller BL, Werck-Reichhart D. Plant cytochrome P450 plasticity and evolution. *Mol Plant* 2021;**14**:1244–65.
14. Riddick DS, Ding X, Wolf CR, Porter TD, Pandey AV, Zhang QY, et al. NADPH-cytochrome P450 oxidoreductase: roles in physiology, pharmacology, and toxicology. *Drug Metab Dispos* 2013;**41**:12–23.
15. Montellano PROD, Vossb JJD. Oxidizing species in the mechanism of cytochrome P450. *Nat Prod Rep* 2002;**19**:477–93.
16. Guengerich FP. Mechanisms of cytochrome P450 substrate oxidation: MiniReview. *J Biochem Mol Toxicol* 2010;**21**:163–8.
17. Ahmed RA, He M, Aftab RA, Zheng S, Nagi M, Bakri R, et al. Bioenergy application of *Dunaliella salina* SA 134 grown at various salinity levels for lipid production. *Sci Rep-UK* 2017;**7**:8118–27.
18. Shi Y, Jiang Z, Hu X, Hu X, Gu R, Jiang B, et al. The cytochrome P450 catalyzing C–S bond formation in *S*-heterocyclization of chuangxinmycin biosynthesis. *Angew Chem Int Ed Engl* 2021;**60**:15399–404.
19. Zhang X, Xu X, You C, Yang C, Guo J, Sang M, et al. Biosynthesis of chuangxinmycin featuring a deubiquitinase-like sulfurtransferase. *Angew Chem Int Ed Engl* 2021;**60**:24418–23.
20. Morita I, Mori T, Mitsuhashi T, Hoshino S, Taniguchi Y, Kikuchi T, et al. Exploiting a C–N bond forming cytochrome P450 monooxygenase for C–S bond formation. *Angew Chem Int Ed Engl* 2020;**59**:3988–93.
21. Grogan G. Hemoprotein catalyzed oxygenations: P450s, UPOs, and progress toward scalable reactions. *JACS Au* 2021;**1**:1312–29.
22. Dixit RB, Suseela MR. Cyanobacteria: potential candidates for drug discovery. *Anton Leeuw Int J G* 2013;**103**:947–61.
23. Harel Y, Ohad I, Kaplan A. Activation of photosynthesis and resistance to photoinhibition in cyanobacteria within biological desert crust. *Plant Physiol* 2004;**136**:3070–9.
24. Radmer RJ. Algal diversity and commercial algal products. *Bioscience* 1996;**46**:263–70.
25. O'Neill E. Mining natural product biosynthesis in eukaryotic algae. *Mar Drugs* 2020;**18**:1–16.
26. Kehr JC, Gatte Picchi D, Dittmann E. Natural product biosyntheses in cyanobacteria: a treasure trove of unique enzymes. *Beilstein J Org Chem* 2011;**7**:1622–35.
27. O'Neill EC, Saalbach G, Field RA. Gene discovery for synthetic biology: exploring the novel natural product biosynthetic capacity of eukaryotic microalgae. *Methods Enzymol* 2016;**576**:99–120.
28. Teng L, Fan X, Nelson DR, Han W, Zhang X, Xu D, et al. Diversity and evolution of cytochromes P450 in stramenopiles. *Planta* 2019;**249**:647–61.
29. Lu J, Shao Y, Qin X, Liu D, Chen A, Li D, et al. CRISPR knockout rat cytochrome P450 3A1/2 model for advancing drug metabolism and pharmacokinetics research. *Sci Rep-UK* 2017;**7**:42922–35.
30. Kim SY, Zhao P, Igarashi M, Sawa R, Tomita T, Nishiyama M, et al. Cloning and heterologous expression of the cyclooctatin biosynthetic gene cluster afford a diterpene cyclase and two p450 hydroxylases. *Chem Biol* 2009;**16**:736–43.
31. Lassen LM, Nielsen AZ, Ziensen B, Gnanasekaran T, Møller BL, Jensen PE. Redirecting photosynthetic electron flow into light-driven synthesis of alternative products including high-value bioactive natural compounds. *ACS Synth Biol* 2014;**3**:1–12.
32. Meunier B, de Visser SP, Shaik S. Mechanism of oxidation reactions catalyzed by cytochrome p450 enzymes. *Chem Rev* 2004;**104**:3947–80.
33. Paine MJ, Scrutton NS, Munro AW, Gutierrez A, Roberts GC, Wolf CR. Electron transfer partners of cytochrome P450. In: *Cytochrome P450-structure, mechanism, and biochemistry*. Boston: Springer; 2005. p. 115–48.
34. Montellano PROD. *Cytochrome P450: structure, mechanism, and biochemistry*. 3rd ed. New York: Springer; 2005.
35. Hausjell J, Halbwirth H, Spadiut O. Recombinant production of eukaryotic cytochrome P450s in microbial cell factories. *Biosci Rep* 2018;**38**:1290–302.
36. Podust LM, Sherman DH. Diversity of P450 enzymes in the biosynthesis of natural products. *Nat Prod Rep* 2012;**29**:1251–66.
37. Jiang Y, Li S. Catalytic function and application of cytochrome P450 enzymes in biosynthesis and organic synthesis. *Chin J Org Chem* 2018;**38**:2307–23.
38. Brash AR. Mechanistic aspects of CYP74 allene oxide synthases and related cytochrome P450 enzymes. *Phytochemistry* 2009;**70**:1522–31.
39. Lawton LA, Codd GA. Cyanobacterial (blue-green-algal) toxins and their significance in UK and European waters. *Water Environ J* 1991;**5**:460–5.
40. Dixit RB, Suseela MR. Cyanobacteria: potential candidates for drug discovery. *Anton Leeuw Int J G* 2013;**103**:947–61.
41. Khumalo MJ, Nzuzi N, Padayachee T, Chen W, Yu JH, Nelson DR, et al. Comprehensive analyses of cytochrome P450 monooxygenases and secondary metabolite biosynthetic gene clusters in cyanobacteria. *Int J Mol Sci* 2020;**21**:656–71.
42. Napoli JL. Physiological insights into all-*trans*-retinoic acid biosynthesis. *Bba-Mol Cell Biol L* 2012;**1821**:152–67.
43. Idres N, Marill J, Flexor MA, Chabot GG. Activation of retinoic acid receptor-dependent transcription by all-*trans*-retinoic acid metabolites and isomers. *J Biol Chem* 2002;**277**:31491–8.
44. Ke N, Baudry J, Makris TM, Schuler MA, Sligar SG. A retinoic acid binding cytochrome P450: CYP120A1 from *Synechocystis* sp. PCC 6803. *Arch Biochem Biophys* 2005;**436**:110–20.
45. Kuhnle K, Ke N, Cryle MJ, Sligar SG, Schuler MA, Schlichting I. Crystal structures of substrate-free and retinoic acid-bound cyanobacterial cytochrome P450 CYP120A1. *Biochemistry* 2008;**47**:6552–9.
46. Alder A, Bigler P, Werck-Reichhart D, Al-Babili S. *In vitro* characterization of *Synechocystis* CYP120A1 revealed the first nonanimal retinoic acid hydroxylase. *FEBS J* 2009;**276**:5416–31.
47. Richard DA, Joseph J, Barchi Jr, Kuniyoshi Masayuki, Moore Richard E, Mynderse JS. Structure of malyngamide C. *J Org Chem* 1985;**50**:2859–62.
48. Moss NA, Leao T, Rankin MR, McCullough TM, Qu P, Korobeynikov A, et al. Ketoreductase domain dysfunction expands chemodiversity: malyngamide biosynthesis in the cyanobacterium *Okeania hirsuta*. *ACS Chem Biol* 2018;**13**:3385–95.
49. Magarvey NA, Beck ZQ, Golakoti T, Ding Y, Huber U, Hemscheidt TK, et al. Biosynthetic characterization and chemoenzymatic assembly of the cryptophycins. Potent anticancer agents from *Nostoc* cyanobionts. *ACS Chem Biol* 2006;**1**:766–79.
50. Smith CD, Zhang XQ, Mooberry SL, Patterson GML, Moore RE. Cryptophycin—a new antimicrotubule agent active against drug-resistant cells. *Cancer Res* 1994;**54**:3779–84.
51. Subbaraju GV, Golakoti T, Patterson GM, Moore RE. Three new cryptophycins from *Nostoc* sp. GSV 224. *J Nat Prod* 1997;**60**:302–5.
52. Ding Y, Seufert WH, Beck ZQ, Sherman DH. Analysis of the cryptophycin P450 epoxidase reveals substrate tolerance and cooperativity. *J Am Chem Soc* 2008;**130**:5492–8.

53. Ramaswamy AV, Sorrels CM, Gerwick WH. Cloning and biochemical characterization of the hectochlorin biosynthetic gene cluster from the marine cyanobacterium *Lyngbya majuscula*. *J Nat Prod* 2007;**70**:1977–86.
54. Liu X, Sheng J, Curtiss 3rd R. Fatty acid production in genetically modified cyanobacteria. *Proc Natl Acad Sci U S A* 2011;**108**:6899–904.
55. Jiang YY, Li Z, Wang C, Zhou YJJ, Xu HF, Li SY. Biochemical characterization of three new alpha-olefin-producing P450 fatty acid decarboxylases with a halophilic property. *Biotechnol Biofuels* 2019;**12**:1–14.
56. Schneider S, Wu Bb, Olts MG, Sanglard D, Witholt B. Biocatalyst engineering by assembly of fatty acid transport and oxidation activities for *in vivo* application of cytochrome P-450BM-3 monooxygenase. *Appl Environ Microbiol* 1998;**64**:3784–90.
57. Lammers PJ, McLaughlin S, Papin S, Trujillo-Provencio C, Ryncarz 2nd AJ. Developmental rearrangement of cyanobacterial *nif* genes: nucleotide sequence, open reading frames, and cytochrome P-450 homology of the *Anabaena* sp. strain PCC 7120 *nifD* element. *J Bacteriol* 1990;**172**:6981–90.
58. Torres S, Fjetland CR, Lammers PJ. Alkane-induced expression, substrate binding profile, and immunolocalization of a cytochrome P450 encoded on the *nifD* excision element of *Anabaena* 7120. *BMC Microbiol* 2005;**5**:5–16.
59. Van den Berg M, Birnbaum L, Bosveld AT, Brunstrom B, Cook P, Feeley M, et al. Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife. *Environ Health Perspect* 1998;**106**:775–92.
60. Duell ER, Milzarek TM, El Omari M, Linares-Otaya LJ, Schäberle TF, König GM, et al. Identification, cloning, expression and functional interrogation of the biosynthetic pathway of the polychlorinated triphenyls ambigol A–C from *Fischerella ambigua* 108b. *Org Chem Front* 2020;**7**:3193–201.
61. Agarwal V, El Gamal AA, Yamanaka K, Poth D, Kersten RD, Schorn M, et al. Biosynthesis of polybrominated aromatic organic compounds by marine bacteria. *Nat Chem Biol* 2014;**10**:640–7.
62. Bouwmeester HJ, Kodde J, Verstappen FW, Altug IG, de Kraker JW, Wallaart TE. Isolation and characterization of two germacrene A synthase cDNA clones from chicory. *Plant Physiol* 2002;**129**:134–44.
63. Agger SA, Lopez-Gallego F, Hoye TR, Schmidt-Dannert C. Identification of sesquiterpene synthases from *Nostoc punctiforme* PCC 73102 and *Nostoc* sp. strain PCC 7120. *J Bacteriol* 2008;**190**:6084–96.
64. Harada H, Shindo K, Iki K, Teraoka A, Okamoto S, Yu F, et al. Efficient functional analysis system for cyanobacterial or plant cytochromes P450 involved in sesquiterpene biosynthesis. *Appl Microbiol Biotechnol* 2011;**90**:467–76.
65. Irie K, Hirota M, Hagiwara N, Koshimizu K, Hayashi H, Murao S, et al. The Epstein-Barr virus early antigen inducing indole alkaloids, (–)-indolactam V and its related compounds, produced by actinomycetes. *Agric Biol Chem* 2014;**48**:1269–74.
66. Huynh MU, Elston MC, Hernandez NM, Ball DB, Kajiyama S, Irie K, et al. Enzymatic production of (–)-indolactam V by LtxB, a cytochrome P450 monooxygenase. *J Nat Prod* 2010;**73**:71–4.
67. Hiraoka M, Shimada S, Uenosono M, Masuda M. A new green-tide-forming alga, *Ulva ohnoi* Hiraoka et Shimada sp. nov. (Ulvales, Ulvophyceae) from Japan. *Phycol Res* 2004;**52**:17–29.
68. Horner RA, Garrison DL, Plumley FG. Harmful algal blooms and red tide problems on the US west coast. *Limnol Oceanogr* 1997;**42**:1076–88.
69. Wang BG, Gloer JB, Ji NY, Zhao JC. Halogenated organic molecules of *Rhodmelaceae* origin: chemistry and biology. *Chem Rev* 2013;**113**:3632–85.
70. Wei X, Hu H, Tong H, Gmitter FG. Profiles of gene family members related to carotenoid accumulation in citrus genus. *J Plant Biol* 2017;**60**:1–10.
71. Gauthier ML, Pickering CR, Miller CJ, Fordyce CA, Chew KL, Berman HK, et al. p38 regulates cyclooxygenase-2 in human mammary epithelial cells and is activated in premalignant tissue. *Cancer Res* 2005;**65**:1792–9.
72. Stumpe M, Feussner I. Formation of oxylipins by CYP74 enzymes. *Phytochemistry Rev* 2006;**5**:347–57.
73. Ohta D, Mizutani M. Redundancy or flexibility: molecular diversity of the electron transfer components for P450 monooxygenases in higher plants. *Front Biosci* 2004;**9**:1587–97.
74. Koeduka T, Ishizaki K, Mwenda CM, Hori K, Sasaki-Sekimoto Y, Ohta H, et al. Biochemical characterization of allene oxide synthases from the liverwort *Marchantia polymorpha* and green microalgae *Klebsormidium flaccidum* provides insight into the evolutionary divergence of the plant CYP74 family. *Planta* 2015;**242**:1175–86.
75. Toporkova YY, Fatykhova VS, Gogolev YV, Khairutdinov BI, Mukhtarova LS, Grechkin AN. Epoxyalcohol synthase of *Ectocarpus siliculosus*. First CYP74-related enzyme of oxylipin biosynthesis in brown algae. *Bba-Mol Cell Biol L* 2017;**1862**:167–75.
76. Ritter A, Dittami SM, Goulitquer S, Correa JA, Boyen C, Potin P, et al. Transcriptomic and metabolomic analysis of copper stress acclimation in *Ectocarpus siliculosus* highlights signaling and tolerance mechanisms in brown algae. *BMC Plant Biol* 2014;**14**:116–32.
77. Dall’Osto L, Fiore A, Cazzaniga S, Giuliano G, Bassi R. Different roles of alpha- and beta-branch xanthophylls in photosystem assembly and photoprotection. *J Biol Chem* 2007;**282**:35056–68.
78. Cui H, Yu X, Wang Y, Cui Y, Li X, Liu Z, et al. Evolutionary origins, molecular cloning and expression of carotenoid hydroxylases in eukaryotic photosynthetic algae. *BMC Genom* 2013;**14**:457–76.
79. Liang MH, Zhu J, Jiang JG. Carotenoids biosynthesis and cleavage related genes from bacteria to plants. *Crit Rev Food Sci Nutr* 2018;**58**:2314–33.
80. Liang MH, Xie H, Chen HH, Liang ZC, Jiang JG. Functional identification of two types of carotene hydroxylases from the green alga *Dunaliella bardawil* rich in lutein. *ACS Synth Biol* 2020;**9**:1246–53.
81. Tamaki S, Kato S, Shinomura T, Ishikawa T, Imaishi H. Physiological role of beta-carotene monohydroxylase (CYP97H1) in carotenoid biosynthesis in *Euglena gracilis*. *Plant Sci* 2019;**278**:80–7.
82. Qi X, Bakht S, Qin B, Leggett M, Hemmings A, Mellon F, et al. A different function for a member of an ancient and highly conserved cytochrome P450 family: from essential sterols to plant defense. *Proc Natl Acad Sci U S A* 2006;**103**:18848–53.
83. Lamb DC, Kelly DE, Kelly SL. Molecular diversity of sterol 14alpha-demethylase substrates in plants, fungi and humans. *FEBS Lett* 1998;**425**:263–5.
84. Lu Y, Zhou W, Wei L, Li J, Jia J, Li F, et al. Regulation of the cholesterol biosynthetic pathway and its integration with fatty acid biosynthesis in the oleaginous microalga *Nannochloropsis oceanica*. *Biotechnol Biofuels* 2014;**7**:81–95.
85. Topletz AR, Thatcher JE, Zelter A, Lutz JD, Tay S, Nelson WL, et al. Comparison of the function and expression of CYP26A1 and CYP26B1, the two retinoic acid hydroxylases. *Biochem Pharmacol* 2012;**83**:149–63.
86. He F, Mori T, Morita I, Nakamura H, Alblova M, Hoshino S, et al. Molecular basis for the P450-catalyzed C–N bond formation in indolactam biosynthesis. *Nat Chem Biol* 2019;**15**:1206–13.
87. Lee DS, Nioche P, Hamberg M, Raman CS. Structural insights into the evolutionary paths of oxylipin biosynthetic enzymes. *Nature* 2008;**455**:363–8.
88. Melis A. Solar energy conversion efficiencies in photosynthesis: minimizing the chlorophyll antennae to maximize efficiency. *Plant Sci* 2009;**177**:272–80.
89. Santos-Merino M, Torrado A, Davis GA, Rottig A, Bibby TS, Kramer DM, et al. Improved photosynthetic capacity and photosystem I oxidation via heterologous metabolite engineering in cyanobacteria. *Proc Natl Acad Sci U S A* 2021;**118**:1–9.

90. Mellor SB, Vinde MH, Nielsen AZ, Hanke GT, Abdiaziz K, Roessler MM, et al. Defining optimal electron transfer partners for light-driven cytochrome P450 reactions. *Metab Eng* 2019;**55**:33–43.
91. Xue Y, Zhang Y, Grace S, He Q. Functional expression of an *Arabidopsis* p450 enzyme, p-coumarate-3-hydroxylase, in the cyanobacterium *Synechocystis* PCC 6803 for the biosynthesis of caffeic acid. *J Appl Phycol* 2013;**26**:219–26.
92. Xue Y, He Q. Cyanobacteria as cell factories to produce plant secondary metabolites. *Front Bioeng Biotechnol* 2015;**3**:57–62.
93. Lassen LM, Nielsen AZ, Olsen CE, Bialek W, Jensen K, Moller BL, et al. Anchoring a plant cytochrome P450 via PsaM to the thylakoids in *Synechococcus* sp. PCC 7002: evidence for light-driven biosynthesis. *PLoS One* 2014;**9**:1–10.
94. Gangl D, Zedler JA, Wlodarczyk A, Jensen PE, Purton S, Robinson C. Expression and membrane-targeting of an active plant cytochrome P450 in the chloroplast of the green alga *Chlamydomonas reinhardtii*. *Phytochemistry* 2015;**110**:22–8.
95. Wlodarczyk A, Gnanasekaran T, Nielsen AZ, Zulu NN, Mellor SB, Luckner M, et al. Metabolic engineering of light-driven cytochrome P450 dependent pathways into *Synechocystis* sp. PCC 6803. *Metab Eng* 2016;**33**:1–11.
96. Hoschek A, Heuschkel I, Schmid A, Buhler B, Karande R, Buhler K. Mixed-species biofilms for high-cell-density application of *Synechocystis* sp. PCC 6803 in capillary reactors for continuous cyclohexane oxidation to cyclohexanol. *Bioresour Technol* 2019;**282**:171–8.
97. Gülin I. Antioxidant activity of caffeic acid (3,4-dihydroxycinnamic acid). *Toxicology* 2006;**217**:213–20.
98. Prasad NR, Karthikeyan A, Karthikeyan S, Reddy BV. Inhibitory effect of caffeic acid on cancer cell proliferation by oxidative mechanism in human HT-1080 fibrosarcoma cell line. *Mol Cell Biochem* 2011;**349**:11–9.
99. Chao PC, Hsu C, Yin MC. Anti-inflammatory and anti-coagulatory activities of caffeic acid and ellagic acid in cardiac tissue of diabetic mice. *Nutr Metab* 2009;**6**:33.
100. Yoshimoto M, Kurata-Azuma R, Fujii M, Hou DX, Ikeda K, Yoshidome T, et al. Enzymatic production of caffeic acid by koji from plant resources containing caffeoylquinic acid derivatives. *Biosci Biotechnol Biochem* 2005;**69**:1777–81.
101. Kim YH, Kwon T, Yang HJ, Kim W, Youn H, Lee JY, et al. Gene engineering, purification, crystallization and preliminary X-ray diffraction of cytochrome P450 p-coumarate-3-hydroxylase (C3H), the *Arabidopsis* membrane protein. *Protein Expr Purif* 2011;**79**:149–55.
102. Sibbesen O, Koch B, Halkier BA, Moller BL. Isolation of the heme-thiolate enzyme cytochrome P-450TYR, which catalyzes the committed step in the biosynthesis of the cyanogenic glucoside dhurrin in *Sorghum bicolor* (L.) Moench. *Proc Natl Acad Sci U S A* 1994;**91**:9740–4.
103. Schuchardt U, Carvalho WA, Spinacé E. Why is it interesting to study cyclohexane oxidation?. *Synlett* 1993:713–8.1993.
104. Schuchardt U, Cardoso D, Sercheli R, Pereira R, Cruz RSd, Guerreiro MC, et al. Cyclohexane oxidation continues to be a challenge. *Appl Catal A-Gen* 2001;**211**:1–17.
105. Salamanca D, Karande R, Schmid A, Dobsław D. Novel cyclohexane monooxygenase from *Acidovorax* sp. CHX100. *Appl Microbiol Biotechnol* 2015;**99**:6889–97.
106. Hoschek A, Toepel J, Hochkeppel A, Karande R, Buhler B, Schmid A. Light-dependent and aeration-independent gram-scale hydroxylation of cyclohexane to cyclohexanol by CYP450 harboring *Synechocystis* sp. PCC 6803. *Biotechnol J* 2019;**14**:724–33.
107. Berepiki A, Hitchcock A, Moore CM, Bibby TS. Tapping the unused potential of photosynthesis with a heterologous electron sink. *ACS Synth Biol* 2016;**5**:1369–75.
108. Kawahigashi H, Hirose S, Ohkawa H, Ohkawa Y. Transgenic rice plants expressing human CYP1A1 remediate the triazine herbicides atrazine and simazine. *J Agric Food Chem* 2005;**53**:8557–64.
109. Inui H, Shiota N, Motoi Y, Ido Y, Inoue T, Kodama T, et al. Metabolism of herbicides and other chemicals in human cytochrome P450 species and in transgenic potato plants co-expressing human CYP1A1, CYP2B6 and CYP2C19. *J Pestic Sci* 2001;**26**:28–40.
110. Mellor SB, Nielsen AZ, Burow M, Motawia MS, Jakubauskas D, Moller BL, et al. Fusion of ferredoxin and cytochrome P450 enables direct light-driven biosynthesis. *ACS Chem Biol* 2016;**11**:1862–9.
111. Kannchen D, Zabret J, Oworah-Nkruma R, Dyczmons-Nowaczyk N, Wiegand K, Lobbert P, et al. Remodeling of photosynthetic electron transport in *Synechocystis* sp. PCC 6803 for future hydrogen production from water. *BBA-Bioenergetics* 2020;**1861**:148208–18.
112. Cui H, Yu X, Wang Y, Cui Y, Li X, Liu Z, et al. Gene cloning and expression profile of a novel carotenoid hydroxylase (CYP97C) from the green alga *Haematococcus pluvialis*. *J Appl Phycol* 2013;**26**:91–103.
113. Yang LE, Huang XQ, Hang Y, Deng YY, Lu QQ, Lu S. The P450-type carotene hydroxylase PuCHY1 from *Porphyra* suggests the evolution of carotenoid metabolism in red algae. *J Integr Plant Biol* 2014;**56**:902–15.
114. Cui H, Ma H, Cui Y, Zhu X, Qin S, Li R. Cloning, identification and functional characterization of two cytochrome P450 carotenoids hydroxylases from the diatom *Phaeodactylum tricorutum*. *J Biosci Biotechnol* 2019;**128**:755–65.